

Limothrips cerealium

Distinguishing features

Female fully winged. Body brown, tarsi and antennal segment III yellowish; fore wings light brown. Antennae 8-segmented; segment III asymmetric, apex with simple sense cone; IV with sense cone simple. Head longer than wide, projecting in front of eyes; three pairs of ocellar setae present, pair III anterolateral to triangle, scarcely longer than distance between two ocelli; maxillary palps 2-segmented. Pronotum with one pair of long posteroangular setae. Metanotum reticulate, campaniform sensilla present, median setae arise behind anterior margin. Fore wing first vein with 2 setae on distal half, second vein with about 8 setae. Abdominal tergites reticulate medially, with one pair of campaniform sensilla close to posterior margin; craspedum not developed; tergite X with one pair of stout thorn-like setae at posterior. Sternites II–VII with 5–10 discal setae, without craspeda.

Male wingless, without ocelli on head; tergite IX medially with pair of stout thorn-like setae on tubercles, posterolateral pair of setae also short and stout; sternites III–VII with small sub-circular pore plate.

Related species

Although eight species are listed in the genus *Limothrips* (eg., zur Strassen, 2003), *L. clarus* was recently declared a *nomen dubium* that was presumably based on a larva, and *L. serotina* was synonymised with *Thrips tabaci* (Mound & Sahin Negus, 2024). The six *Limothrips* species all live on various grasses, and are distinguished from other grass-living Thripinae by the stout pair of thorn-like setae at the apex of the tenth abdominal segment. Although they are originally from Europe, three species are now widespread around the world. *L. cerealium* has neither antennal segment III produced outwards nor stout setae on tergite IX.

Biological data

Feeding and breeding on the leaves and in the flowers of grasses and cereal crops. Following the build-up of large populations in summer cereal crops this thrips can appear in swarms, and is referred to colloquially as thunderflies. Kirk (2004) discussed possible causal mechanisms for this phenomenon that is commonly associated with stormy weather in Britain and other parts of Europe.

Distribution data

Widespread and common, and recorded across Britain from Kent to the Orkney Islands, and also from both Northern Ireland and the Republic of Ireland (Mound *et al.*, 1976; O'Connor, 2008), this species is abundant across Europe. It has also been introduced into, and is now found in, many temperate areas around the world.

Family name

THRIPIDAE - THRIPINAE

Species name



Female



Male



Head & pronotum



Head & pronotum

Limothrips cerealium (Haliday)

Original name and synonyms

Thrips (*Limothrips*) *cerealium* Haliday, 1836: 445
Limothrips avenae Hinds, 1902: 139
Limothrips cerealium f. *aptera* Karny, 1914: 56
Limothrips minor Bagnall, 1927: 565
Limothrips cerealium f. *adusta* Maltbaek, 1927: 14
Limothrips cerealium ssp. *astutus* Priesner, 1964: 115
Limothrips syriacus Jenser, 2009: 83

References

Kirk WDJ (2004) The link between cereal thrips and thunderstorms. *Acta Phytopathologica et Entomologica Hungarica* **39** (1-3): 131–136.

Mound LA, Morison GD, Pitkin BR & Palmer JM (1976) Thysanoptera. *Handbooks for the Identification of British Insects* **1** (11): 1–79.

Mound, L.A. & Şahin Negiş, İ. (2024). Two enigmatic species epithets in the genus *Limothrips* (Thysanoptera, Thripinae). *Zootaxa*, 5512 (4), 597–598.

O'Connor JP (2008) A review of the Irish thrips (Thysanoptera). *Irish Naturalists' Journal* **29**: 20–24.

zur Strassen R (2003) Die terebranten Thysanopteren Europas und des Mittelmeer-Gebietes. *Die Tierwelt Deutschlands* **74**: 1–271.




Antenna




Antenna



Meso & metanota




Female tergites VIII-X




Thoracic sternites




Fore wing



Male tergite IX



Male tergite IX



Male pore plates