

Apterothrips apteris

Distinguishing features

Both sexes without wings, and uniformly dark brown. Antennae 8-segmented but VI with an incomplete oblique suture producing an apparently 9-segmented condition; segments III–IV with sense cone simple. Head weakly reticulate; ocelli not developed, ocellar setae III as long as diameter of three ommatidia; maxillary palps 3-segmented. Pronotum with no long setae. Tergites with transverse reticulation medially, tergites I–VIII with well-developed craspedum on posterior margin. Sternites III–VI with well-developed 5-lobed craspedum, the sternal posteromarginal setal pair S3 arising at the posterolateral angle of each of the sternites. Male smaller and paler than female, sternite II with an internal glandular structure opening through an aperture on the antecostal margin of segment III; tergite IX with 2 pairs of short stout thorn-like setae.

Related species

There are only two species of *Apterothrips*, of which *A. apteris* is essentially a southern hemisphere species whereas *A. secticornis* is essentially a northern hemisphere species (Mound & Marullo, 1996). These two are distinguished by the number of lobes on the craspeda of the abdominal sternites, but are otherwise very similar. The males share with those of several Neotropical Thripidae species the character state of having a single glandular area on the anterior margin of the third abdominal sternite.

Biological data

Feeding and breeding on the leaves of its host plants, and usually associated with Poaceae. However, this thrips has been found causing damage to garlic and to lucerne in Australia, and large populations have been recorded in California from *Erigeron* [Asteraceae].

Distribution data

In Britain, recorded twice in Scotland, at Ayr and Fair Isle (Morison, 1973a), and twice in England, in West Yorkshire and Devon (Collins, 2000a; 2010a), but otherwise not known from Europe. The species is widespread around the southern hemisphere from Mexico, Peru, and the Falkland Islands, to Australia and New Zealand, and is also known from Hawaii. The British records are therefore geographically aberrant. It was previously misidentified as *secticornis*, a species that, by contrast, is reported across Europe and the northern Holarctic, but is not known from Britain.

Family name

THRIPIDAE - THRIPINAE


Species name

Apterothrips apteris (Daniel)

Original name and synonyms

- Sericothrips apteris* Daniel, 1904: 295
- Sericothrips stanfordii* Moulton, 1907: 51
- Sericothrips ineptus* Ahlberg, 1922: 271
- Apterothrips delamarei* Bournier, 1962: 231

References




Collins DW (2000a) First English record for *Apterothrips apteris* (Daniel) (Thysanoptera: Thripidae). *Entomologist's Monthly Magazine*. **136**: 210.

Collins DW (2010a) Thysanoptera of Great Britain: a revised and updated checklist. *Zootaxa* **2412**: 21–41.

Morison GD (1973a) Observations and records for British Thysanoptera. XVI. Thripidae, *Apterothrips secticornis* (Trybom), a species new to Britain. *Entomologist's Monthly Magazine* **109**: 208–210.

Mound LA & Marullo R (1996) The thrips of Central and South America: an introduction (Insecta: Thysanoptera). *Memoirs on Entomology, International* **6**: 1–487.



Male glandular opening between sternites II and III