

# Edgbastonia (Barcaldinia) hufferensis Ponder, Zhang, Hallan & Shea, 2019



The state of the s



Distribution of Edgbastonia (Barcaldinia) hufferensis

Spring at Lake Huffer, the habitat of this species.

Photo A. Davis.

Edgbastonia (Barcaldinia) hufferensis (adult size 1.7 - 2.2 mm)

## **Diagnostic features**

The most similar species is *E.* (*B.*) *colmani*, but it differs in molecular results and in the male having an almost straight pallial vas deferens and penial duct, although the female reproductive system is similar to that of *E.* (*B.*) *colmani*. Compared with *E.* (*B.*) *pallida* this species is larger and darker coloured.

#### Classification

Edgbastonia (Barcaldinia) hufferensis Ponder, Zhang, Hallan & Shea, 2019

Class Gastropoda

Infraclass Caenogastropoda

Order Littorinida

Suborder Rissoidina

Superfamily Truncatelloidea

Family Tateidae

Genus Edabastonia Ponder in Ponder, Wilke, Zhang, Golding, Fukuda, & Mason 2008 (Type species: Edgbastonia alanwillsi Ponder in Ponder et al., 2008).

Subgenus Barcaldinia Ponder, Zhang, Hallan & Shea, 2019 (Type species Jardinella edgbastonensis Ponder & Clark, 1990)

Original name: Edgbastonia (Barcaldinia) hufferensis Ponder, Zhang, Hallan & Shea, 2019. In Ponder, W. F., Zhang, W.-H., Hallan, A., & Shea, M. E. (2019). New taxa of Tateidae (Caenogastropoda, Truncatelloidea) from springs associated with the Great Artesian Basin and Einasleigh Uplands, Queensland, with the description of two related taxa from eastern coastal drainages. Zootaxa 4583(1): 1-67.

Type locality: Near base of main spring, Lake Huffer Station, approx. 0.3 km from spring head, Qeensland.

### Biology and ecology

Lives on sandy substrate and vegetation in the spring.

#### **Distribution**

Main spring on Lake Huffer Station, north of Aramac, Queensland (Barcaldine Supergroup). This spring is approximately seven km SSW of the type locality of E. (B.) colmani.

## **Further reading**

Fensham, R., Ponder, W. & Fairfax, R. (2010). Recovery plan for the community of native species dependent on natural discharge of groundwater from the Great Artesian Basin. Report to Department of the Environment, Water, Heritage and the Arts, Canberra. Queensland Department of Environment and Resource Management, Brisbane. https://www.environment.gov.au/system/files/resources/0cefc83a-3854-4cff-9128-abc719d9f9b3/files/great-artesian-basin-ec.pdf

Ponder, W. F., Zhang, W. -H., Hallan, A., & Shea, M. E. (2019). New taxa of Tateidae (Caenogastropoda, Truncatelloidea) from springs associated with the Great Artesian Basin and Einasleigh Uplands, Queensland, with the description of two related taxa from eastern coastal drainages. Zootaxa 4583(1): 1-67.

To cite this resource: Ponder, W. F., Hallan, A., Shea, M. E., Clark, S. A., Richards, K., Klunzinger, M. W., and Kessner, V. 2023. Australian Freshwater Molluscs. Revision 2.

https://keys.lucidcentral.org/keys/v3/freshwater\_molluscs/

To contact the authors for comment or suggestions, please email: fwmollusc@gmail.com

Copyright © 2023. All rights reserved. The Australian Museum.









