



***Hemistomia beaumonti* Ponder, 1982**

Diagnostic features

This species, together with *H. flexicolumella*, differs from others on



Hemistomia beaumonti (adult size 2.6-2.9 mm)



Distribution of *Hemistomia beaumonti* (map showing Lord Howe Island, Australia).

Lord Howe Island in having a columellar bulge and an operculum with a rudimentary peg and no white smear or spot. The shell is sculptured with growth lines only and the suture is simple or weakly channelled. The maximum length of the shell is about 2.9 mm

The penis is simple in all *Hemistomia* species.

Classification

Hemistomia beaumonti Ponder, 1982

Class Gastropoda

Infraclass Caenogastropoda

Order Littorinida

Suborder Rissoidina

Superfamily Truncatelloidea

Family Tateidae

Genus Hemistomia Crosse, 1872

Original name: Hemistomia beaumonti Ponder, 1982. In Ponder, W.F. (1982) Hydrobiidae of Lord Howe Island (Mollusca: Gastropoda: Prosobranchia). *Australian Journal of Marine and Freshwater Research* 33: 89-159.

Type locality: Southern tributary of southern end of main gully in middle of Mt. Gower, Lord Howe Island.

Biology and ecology

Found in streams, seepages and pools. It is assumed that like most tateids, the Lord Howe Island Tateidae feed on bacteria, microscopic algae, diatoms and possibly decaying vegetation.

Distribution

Restricted to the top of Mt Gower, Lord Howe Island.

Further reading

Ponder, W. F. (1982). Hydrobiidae of Lord Howe Island (Mollusca: Gastropoda: Prosobranchia). *Australian Journal of Marine and Freshwater Research* 33: 89-159.

To cite this resource: **Ponder, W. F., Hallan, A., Shea, M. E., Clark, S. A., Richards, K., Klunzinger, M. W., and Kessner, V. 2023. Australian Freshwater Molluscs. Revision 2.**

https://keys.lucidcentral.org/keys/v3/freshwater_molluscs/

To contact the authors for comment or suggestions, please email: fwmollusc@gmail.com

Copyright © 2023. All rights reserved. The Australian Museum.

