


Aeolothrips fasciatus



Distinguishing features

Female macropterous. Body and legs brown; antennal segment III yellow with extreme apex brown, IV–IX brown; fore wing dark at extreme base but clavus largely pale, with two transverse dark bands, median pale band shorter than distal dark band, apex clear with ring vein pale. Head with postocular region as long as eye length; maxillary palp 3-segmented. Antennae 9-segmented; segment III long and slender with linear sensorium scarcely 0.3 as long as segment, IV with sensorium almost 0.5 as long as segment and curved distally, without internal markings. Mesonotum with no accessory setae medially. Metanotal reticles without internal markings. Abdominal tergite I with faint transverse lines anteromedially; trichobothria on X no larger than base of major setae on X. Sternites with 4 pairs of marginal setae, discal setae not developed; sternite VII accessory setae arising well in front of margin.

Male similar to female but smaller; antennal segment III almost fuscous rather than yellow, but abdominal segments III–V sometimes paler than II and VI. Abdominal tergite I with two longitudinal ridges; IV–V each with pair of irregular tubercles; IX with paired bifurcate claspers postero-laterally, posteroangular pair of setae longer than claspers. Sternites without discal setae.



Related species

The genus *Aeolothrips* is essentially Holarctic, and currently includes about 90 species. However, *A. fasciatus* can be recognised unequivocally only in the male sex, in which the posteroangular setae on tergite IX are longer than the claspers. Females are closely similar to *A. intermedius*, a very common species in Europe. However, *A. intermedius* has the third antennal segment less slender than in *A. fasciatus* and more extensively brown toward the apex. *A. fasciati pennis* Blanchard from Chile is probably a parthenogenetic strain of *A. fasciatus*, and apparently identical females are known also from California. Based on males and females, *A. fasciatus* is recorded from Tasmania, and also from New Zealand. It is readily distinguished from *Desmothrips* species by the very different banding pattern on the fore wings with the wing apex pale.

Biological data

Breeding in the flowers of plants in various families, with no recorded specificity, but presumably also acting as a facultative predator of other thrips

Distribution data

Described from Europe, this species has been recorded in various parts of the world. However, records based only on females may not be correct, because only in the male sex can *A. fasciatus* be distinguished unequivocally. Males of this species have been studied from Finland, Denmark, Germany, Slovakia, Azores, Hokkaido, New Zealand and Tasmania.

Family name

AEOLOTHRIPIDAE

Species name

Aeolothrips fasciatus (Linnaeus)

Original name and synonyms

Thrips fasciata Linnaeus, 1758: 457

Coleothrips trifasciatus Fitch, 1855: 386

Aeolothrips fasciatus f. *atricornis* Maltbaek, 1927: 14

Aeolothrips fasciatus justiciae Post, 1961: 137

References

Mound L, Collins D & Hastings A (201) *Thysanoptera Britannica et Hibernica. A guide to British thrips*. Lucidcentral.org, Identic Pty Ltd, Queensland, Australia. https://keys.lucidcentral.org/keys/v3/british_thrips/

ThripsWiki (2020) *ThripsWiki - providing information on the World's thrips*. Available from: http://thrips.info/wiki/Main_Page [accessed 12.xii.2019].