


# Thrips roepkei


## Distinguishing features

Both sexes fully winged. Body light brown, femora brown to yellowish brown, tibiae and tarsi often almost yellow; major setae light brown; antennal segment III almost yellow, IV–V yellow but light brown toward apex, VI–VII brown; fore wings pale to scarcely shaded. Antennae 7-segmented; III & IV each with short forked sense cone, VII small. Head with 2 pairs of ocellar setae; pair III no longer than distance between 2 ocelli, arising on anterior margins of triangle; postocular setal pairs I & III longer than ocellar setae III, postocular setal pair II small. Pronotum with 2 pairs of long posteroangular setae; posterior margin with 3 pairs of setae; discal area with widely spaced transverse striae. Mesonotum with paired anterior campaniform sensilla; median setae arise well in front of posterior margin. Metanotum with irregular reticulation medially, elongate on posterior third; median setae arising behind anterior margin; campaniform sensilla absent. Fore wing first vein with 3 setae on distal half; second vein with 11–12 setae. Abdominal tergite II with 4 lateral marginal setae; tergites V–VIII with paired ctenidia, on VIII posteromesad to spiracles; tergite VIII posteromarginal comb absent medially, with few or no microtrichia laterally; pleurotergites with no discal setae, sculpture lines with a few dentate microtrichia, posterior margin with few or no microtrichia; tergite IX with 2 pairs of campaniform sensilla, X with median split. Sternites with no discal setae; sternite VII marginal setae S1 arise close to posterior margin. Male yellow, smaller than female; tergite VIII with no posteromarginal comb; tergite IX median setae slender; sternites III–VII with transverse pore plate.



Head

Pronotum



Meso & metanota

Meso & metanota

Tergites I-II



Sternites VI-VII

## Related species

This monophagous species is similar in structure to the widespread polyphagous species *Thrips fuscipennis*, but the fore wings are much paler, and the tibiae as well as the tarsi are almost yellow. Vierbergen (2004) synonymised *fallaciosus* with *roepkei*, but the North American species is reported to have a wider host range, and the relationship between these two needs further investigation. The genus *Thrips* is the second largest genus in the Thysanoptera, and currently includes, worldwide, over 290 species. All members of genus *Thrips* lack ocellar setae I on the head, and they all have ctenidia on tergite VIII posteromesad to the spiracles. Other characters, such as number of antennal segments, number of setae on the fore wing veins, and number of discal setae on the sternites are variable between species (Palmer, 1992; Nakahara, 1994; Mound & Masumoto, 2005).

## Biological data

Feeding and breeding in the flowers of its host plants, *Solanum dulcamara* [Solanaceae]. The initial report of *Solanum nigrum* as the host may be the result of erroneous plant species identification (Vierbergen, 2004).

## Distribution data

In Europe, recorded from southern Britain, Norway, the Netherlands and Germany. Also widespread in northern North America, assuming that the synonymy of *fallaciosus* is correct.

## Family name

THRIPIDAE - THRIPINAE

## Species name

*Thrips roepkei* Doeksen

## Original name and synonyms

*Thrips roepkei* Doeksen, 1953: 169  
*Thrips inopinatus* zur Strassen, 1963: 523  
*Thrips fallaciosus* Nakahara, 1994: 51.

## References

Mound LA & Masumoto M (2005) The genus *Thrips* (Thysanoptera, Thripidae) in Australia, New Caledonia and New Zealand. *Zootaxa* **1020**: 1–64.

Nakahara S (1994) The genus *Thrips* Linnaeus (Thysanoptera: Thripidae) of the New World. *United States Department of Agriculture. Technical Bulletin* **1822**: 1–183.

Palmer JM (1992) *Thrips* (Thysanoptera) from Pakistan to the Pacific: a review. *Bulletin of the British Museum (Natural History) Entomology Series* **61** (1): 1–76.

Vierbergen G (2004) Eight species of thrips new for the Netherlands and some taxonomical changes in *Stenchaetothrips*, *Thrips* and *Hoplothrips*. *Acta Phytopathologica et Entomologica Hungarica* **39**: 199–209.