

# Pterostylis atrosanguinea

## Name and Authority

**Pterostylis atrosanguinea** (D.L.Jones & C.J.French) D.L.Jones & C.J.French

## Synonyms (previous names)

*Pterostylis* sp. 'crowded'; *Urochilus atrosanguineus*

## Common Name

Crowded Banded Greenhood

## Complex Name

*Pterostylis vittata* (Banded Greenhoods)

## Originally Described by

David Jones & Christopher French in *Australian Orchid Review* 82(3): 49-51, Fig. 1 (2017) as *Urochilus atrosanguineus*

## Distribution and Habitat

Toodyay to Bunbury and eastwards to Kellerberrin, growing in sandy and sandy-clay soils in forests, woodlands and shrublands.

## Flowering Months

· · · · · Jul Aug Sep · · ·

## Description

A robust **species** 200–400 mm high with 10 to 20 **cauline** leaves 40–60 mm long by 4–14 mm wide and up to 20 semi translucent, dark reddish-brown and white flowers 14–20 mm across.

## Distinguishing Features

It is distinguished from the related Dark Banded Greenhood (*Pterostylis sanguinea*) by its more robust appearance and larger, more numerous, darker coloured flowers.

## WA Conservation Code (Threatened Status)

Not Listed

Copyright © 2024. WANOSCG All rights reserved. (ver: April 2024)



*Pterostylis atrosanguinea*



*Pterostylis atrosanguinea* distribution



*Pterostylis atrosanguinea*



*Pterostylis atrosanguinea*



*Pterostylis atrosanguinea*

