

# Pterostylis concava

## Name and Authority

**Pterostylis concava** D.L.Jones & M.A.Clem.

## Synonyms (previous names)

*Pterostylis vittata* var. *subdifformis*; *Oligochaetochilus concavus*; *Urochilus concavus*

## Common Name

Cupped Banded Greenhood

## Complex Name

*Pterostylis vittata* (Banded Greenhoods)

## Originally Described by

David Jones & Mark Clements in *Austral.Orchid Res.* 1:121 (1989)

## Distribution and Habitat

Calingiri to Hopetoun, growing in sandy, sandy-clay and lateritic soils in forests, woodlands and shrublands.

## Flowering Months

· · · · · Jun Jul Aug · · · · ·

## Description

A distinctive [species](#) 100–350 mm high with three to nine [cauline](#) leaves 14–40 mm long by 4–7 mm wide and up to 10 semi translucent, reddish-brown and white flowers 12–18 mm across.

## Distinguishing Features

It is distinguished from the other members of the *Pterostylis vittata* [complex](#) by its prominently cupped [lateral sepals](#).

## WA Conservation Code (Threatened Status)

Not Listed



[Pterostylis concava](#)



[Pterostylis concava distribution](#)



[Pterostylis concava](#)



[Pterostylis concava](#)



[Pterostylis concava](#)

Copyright © 2024. WANOSCG All rights reserved. (ver: April 2024)



Web edition hosted at <https://keys.lucidcentral.org/keys/v4/wa-orchids/>